


Amendments to the Claims

This listing of claims will replace all prior versions, and listings, of claims in the application:

Claim 1 (Currently amended): A compound of formula III:



or a pharmaceutically acceptable derivative or prodrug salt thereof, wherein:

Z¹ is nitrogen or CR⁸, Z² is CH, and Z³ is nitrogen or CR⁸, provided that when one of Z¹ or Z³ is nitrogen, the other of Z¹ or Z³ is CR⁸ or CR⁸, respectively;

R⁸ is T-R³ or L-Z-R³;

Q is selected from -N(R⁴)-, -O-, -S-, or -CH(R⁶)-;

R¹ is T-(Ring D);

Ring D is a 5-7 membered monocyclic ring or 8-10 membered bicyclic ring selected from aryl, heteroaryl, heterocyclyl or carbocyclyl, said heteroaryl or heterocyclyl ring having 1-4 ring heteroatoms selected from nitrogen, oxygen or sulfur, wherein each substitutable ring carbon of Ring D is independently substituted by oxo, T-R⁵, or V-Z-R⁵, and each substitutable ring nitrogen of Ring D is independently substituted by -R⁴;

T is a valence bond or a C₁₋₄ alkylidene chain, wherein when Q is -CH(R⁶)-, a methylene unit of said C₁₋₄ alkylidene chain is optionally replaced by -O-, -S-, -N(R⁴)-, -CO-, -OC(O)NH-, or -NHCO₂-;

Z is a C₁₋₄ alkylidene chain;

L is -O-, -S-, -SO-, -SO₂-, -N(R⁶)SO₂-, -SO₂N(R⁶)-, -N(R⁶)-, -CO-, -CO₂-, -N(R⁶)CO-, -N(R⁶)C(O)O-, -N(R⁶)CON(R⁶)-, -N(R⁶)SO₂N(R⁶)-, -N(R⁶)N(R⁶)-, -C(O)N(R⁶)-, -OC(O)N(R⁶)-, -C(R⁶)₂O-, -C(R⁶)₂S-, -C(R⁶)₂SO-, -C(R⁶)₂SO₂-, -C(R⁶)₂SO₂N(R⁶)-,

$-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$, $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, or $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$;

R^2 and $\text{R}^{2'}$ are independently selected from $-\text{R}$, $-\text{T}-\text{W}-\text{R}^6$, or R^2 and $\text{R}^{2'}$ are taken together with their intervening atoms to form a fused, 5-8 membered, unsaturated or partially unsaturated, ring having 0-3 ring heteroatoms selected from nitrogen, oxygen, or sulfur, wherein each substitutable ring carbon of said fused ring formed by R^2 and $\text{R}^{2'}$ is independently substituted by halo, oxo, $-\text{CN}$, $-\text{NO}_2$, $-\text{R}^7$, or $-\text{V}-\text{R}^6$, and each substitutable ring nitrogen of said ring formed by R^2 and $\text{R}^{2'}$ is independently substituted by R^4 ;

R^3 is selected from $-\text{R}$, $-\text{halo}$, $-\text{OR}$, $-\text{C}(=\text{O})\text{R}$, $-\text{CO}_2\text{R}$, $-\text{COCOR}$, $-\text{COCH}_2\text{COR}$, $-\text{NO}_2$, $-\text{CN}$,
 $-\text{S}(\text{O})\text{R}$, $-\text{S}(\text{O})_2\text{R}$, $-\text{SR}$, $-\text{N}(\text{R}^4)_2$, $-\text{CON}(\text{R}^7)_2$, $-\text{SO}_2\text{N}(\text{R}^7)_2$, $-\text{OC}(=\text{O})\text{R}$, $-\text{N}(\text{R}^7)\text{COR}$,
 $-\text{N}(\text{R}^7)\text{CO}_2(\text{C}_{1-6} \text{ aliphatic})$, $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$, $-\text{C}=\text{NN}(\text{R}^4)_2$, $-\text{C}=\text{N}-\text{OR}$, $-\text{N}(\text{R}^7)\text{CON}(\text{R}^7)_2$,
 $-\text{N}(\text{R}^7)\text{SO}_2\text{N}(\text{R}^7)_2$, $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^7)_2$;

each R is independently selected from hydrogen or an optionally substituted group selected from C_{1-6} aliphatic, C_{6-10} aryl, a heteroaryl ring having 5-10 ring atoms, or a heterocyclyl ring having 5-10 ring atoms;

each R^4 is independently selected from $-\text{R}^7$, $-\text{COR}^7$, $-\text{CO}_2(\text{optionally substituted } \text{C}_{1-6} \text{ aliphatic})$, $-\text{CON}(\text{R}^7)_2$, or $-\text{SO}_2\text{R}^7$;

each R^5 is independently selected from $-\text{R}$, halo, $-\text{OR}$, $-\text{C}(=\text{O})\text{R}$, $-\text{CO}_2\text{R}$, $-\text{COCOR}$, $-\text{NO}_2$, $-\text{CN}$,
 $-\text{S}(\text{O})\text{R}$, $-\text{SO}_2\text{R}$, $-\text{SR}$, $-\text{N}(\text{R}^4)_2$, $-\text{CON}(\text{R}^4)_2$, $-\text{SO}_2\text{N}(\text{R}^4)_2$, $-\text{OC}(=\text{O})\text{R}$, $-\text{N}(\text{R}^4)\text{COR}$,
 $-\text{N}(\text{R}^4)\text{CO}_2(\text{optionally substituted } \text{C}_{1-6} \text{ aliphatic})$, $-\text{N}(\text{R}^4)\text{N}(\text{R}^4)_2$, $-\text{C}=\text{NN}(\text{R}^4)_2$, $-\text{C}=\text{N}-\text{OR}$,
 $-\text{N}(\text{R}^4)\text{CON}(\text{R}^4)_2$, $-\text{N}(\text{R}^4)\text{SO}_2\text{N}(\text{R}^4)_2$, $-\text{N}(\text{R}^4)\text{SO}_2\text{R}$, or $-\text{OC}(=\text{O})\text{N}(\text{R}^4)_2$;

V is $-\text{O}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{N}(\text{R}^6)\text{SO}_2-$, $-\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{N}(\text{R}^6)-$, $-\text{CO}-$, $-\text{CO}_2-$, $-\text{N}(\text{R}^6)\text{CO}-$,
 $-\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$, $-\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{O})\text{N}(\text{R}^6)-$,
 $-\text{OC}(\text{O})\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{O}-$, $-\text{C}(\text{R}^6)_2\text{S}-$, $-\text{C}(\text{R}^6)_2\text{SO}-$, $-\text{C}(\text{R}^6)_2\text{SO}_2-$, $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$, $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$, $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$, or $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$;


W is $-\text{C}(\text{R}^6)_2\text{O}-$, $-\text{C}(\text{R}^6)_2\text{S}-$, $-\text{C}(\text{R}^6)_2\text{SO}-$, $-\text{C}(\text{R}^6)_2\text{SO}_2-$, $-\text{C}(\text{R}^6)_2\text{SO}_2\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)-$, $-\text{CO}-$,
 $-\text{CO}_2-$, $-\text{C}(\text{R}^6)\text{OC}(\text{O})-$, $-\text{C}(\text{R}^6)\text{OC}(\text{O})\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CO}-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{C}(\text{O})\text{O}-$,
 $-\text{C}(\text{R}^6)=\text{NN}(\text{R}^6)-$, $-\text{C}(\text{R}^6)=\text{N}-\text{O}-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{N}(\text{R}^6)-$, $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{SO}_2\text{N}(\text{R}^6)-$,
 $-\text{C}(\text{R}^6)_2\text{N}(\text{R}^6)\text{CON}(\text{R}^6)-$, or $-\text{CON}(\text{R}^6)-$;

each R^6 is independently selected from hydrogen or an optionally substituted C_{1-4} aliphatic group, or two R^6 groups on the same nitrogen atom are taken together with the nitrogen atom to form a 5-6 membered heterocyclyl or heteroaryl ring;

each R^7 is independently selected from hydrogen or an optionally substituted C_{1-6} aliphatic group, or two R^7 on the same nitrogen are taken together with the nitrogen to form a 5-8 membered heterocyclyl or heteroaryl ring; and

R^8 is selected from -R, halo, -OR, -C(=O)R, -CO₂R, -COCOR, -NO₂, -CN, -S(O)R, -SO₂R, -SR, -N(R⁴)₂, -CON(R⁴)₂, -SO₂N(R⁴)₂, -OC(=O)R, -N(R⁴)COR, -N(R⁴)CO₂(optionally substituted C_{1-6} aliphatic), -N(R⁴)N(R⁴)₂, -C=NN(R⁴)₂, -C=N-OR, -N(R⁴)CON(R⁴)₂, -N(R⁴)SO₂N(R⁴)₂, -N(R⁴)SO₂R, or -OC(=O)N(R⁴)₂.

Claim 2 (Currently amended): The compound according to claim 1, wherein Q is -N(R⁴)-, -S-, or -CH(R⁶)-, and said compound is of formula IIIa or IIIb



or a pharmaceutically acceptable derivative or prodrug salt thereof.

Claim 3 (Original): The compound according to claim 2, wherein said compound has one or more features selected from the group consisting of:

- (a) R^x is hydrogen, alkyl- or dialkylamino, acetamido, or a C_{1-4} aliphatic group;
- (b) R^1 is T-(Ring D), wherein T is a valence bond or a methylene unit;
- (c) Ring D is a 5-7 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (d) R^2 is -R or -T-W- R^6 and $R^{2'}$ is hydrogen, or R^2 and $R^{2'}$ are taken together to form an optionally substituted benzo ring.

Claim 4 (Original): The compound according to claim 3, wherein:

- (a) R^x is hydrogen, alkyl- or dialkylamino, acetamido, or a C_{1-4} aliphatic group;
- (b) R^1 is T-(Ring D), wherein T is a valence bond or a methylene unit;
- (c) Ring D is a 5-7 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (d) R^2 is $-R$ or $-T-W-R^6$ and $R^{2'}$ is hydrogen, or R^2 and $R^{2'}$ are taken together to form an optionally substituted benzo ring.

Claim 5 (Original): The compound according to claim 3, wherein said compound has one or more features selected from the group consisting of:

- (a) R^1 is T-(Ring D), wherein T is a valence bond, and Q is $-S-$ or $-NH-$;
- (b) Ring D is a 5-6 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (c) R^2 is $-R$ and $R^{2'}$ is hydrogen, wherein R is selected from hydrogen, C_{1-6} aliphatic, phenyl, a 5-6 membered heteroaryl ring, or a 5-6 membered heterocyclic ring.

Claim 6 (Original): The compound according to claim 5, wherein:

- (a) R^1 is T-(Ring D), wherein T is a valence bond, and Q is $-S-$ or $-NH-$;
- (b) Ring D is a 5-6 membered monocyclic or an 8-10 membered bicyclic aryl or heteroaryl ring; and
- (c) R^2 is $-R$ and $R^{2'}$ is hydrogen, wherein R is selected from hydrogen, C_{1-6} aliphatic, phenyl, a 5-6 membered heteroaryl ring, or a 5-6 membered heterocyclic ring.

Claim 7 (Original): The compound according to claim 5, wherein said compound has one or more features selected from the group consisting of:

- (a) R^x is hydrogen methyl, ethyl, propyl, cyclopropyl, isopropyl, methylamino or acetamido;
- (b) R^1 is T-(Ring D), wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from $-halo$, $-CN$, $-NO_2$, $-N(R^4)_2$, optionally substituted C_{1-6} aliphatic group,

- OR, -CO₂R, -CON(R⁴)₂, -OCO(R⁴)₂, -N(R⁴)COR, -N(R⁴)SO₂R,
 -N(R⁶)COCH₂CH₂N(R⁴)₂, or -N(R⁶)COCH₂CH₂CH₂N(R⁴)₂; and
 (c) R² is hydrogen or a substituted or unsubstituted C₁₋₆ aliphatic.

Claim 8 (Original): The compound according to claim 7, wherein:

- (a) R^x is hydrogen methyl, ethyl, propyl, cyclopropyl, isopropyl, methylamino or acetamido;
 (b) R¹ is T-(Ring D), wherein T is a valence bond and Ring D is a 5-6 membered aryl or heteroaryl ring, wherein Ring D is optionally substituted with one to two groups selected from -halo, -CN, -NO₂, -N(R⁴)₂, optionally substituted C₁₋₆ aliphatic group, -OR, -CO₂R, -CON(R⁴)₂, -OCO(R⁴)₂, -N(R⁴)COR, -N(R⁴)SO₂R, -N(R⁶)COCH₂CH₂N(R⁴)₂, or -N(R⁶)COCH₂CH₂CH₂N(R⁴)₂; and
 (c) R² is hydrogen or a substituted or unsubstituted C₁₋₆ aliphatic.

Claim 9 (Previously presented): A compound selected from the group consisting of:

- N⁵-(1*H*-Indazol-6-yl)-N³-(5-methyl-1*H*-pyrazol-3-yl)-pyridazine-3,5-diamine;
 N-{4-[6-(5-Methyl-1*H*-pyrazol-3-ylamino)-pyridazin-4-ylsulfanyl]-phenyl}-acetamide;
 [5-(3-Methoxy-benzyl)-pyridazin-3-yl]-(5-methyl-1*H*-pyrazol-3-yl)-amine;
 N³-(5-Cyclopropyl-1*H*-pyrazol-3-yl)-N⁵-pyridin-3-ylmethyl-pyridazine-3,5-diamine;
 [5-(Benzothiazol-6-ylsulfanyl)-pyridazin-3-yl]-(5-cyclopropyl-1*H*-pyrazol-3-yl)-amine;
 {4-[6-(5-Cyclopropyl-1*H*-pyrazol-3-ylamino)-pyridazin-4-yloxy]-phenyl}-acetonitrile;
 N-{4-[6-(1*H*-Indazol-3-ylamino)-pyridazin-4-ylamino]-phenyl}-methanesulfonamide;
 (1*H*-Indazol-3-yl)-[5-(thiophen-2-ylmethylsulfanyl)-pyridazin-3-yl]-amine;
 N⁵-(5-Methyl-1*H*-pyrazol-3-yl)-N³-pyridin-3-ylmethyl-pyridazine-3,5-diamine;
 [6-(Benzothiazol-6-ylsulfanyl)-pyridazin-4-yl]-(5-methyl-1*H*-pyrazol-3-yl)-amine;
 {4-[5-(5-Methyl-1*H*-pyrazol-3-ylamino)-pyridazin-3-yloxy]-phenyl}-acetonitrile;
 N⁵-(5-Cyclopropyl-1*H*-pyrazol-3-yl)-N³-(1*H*-indazol-6-yl)-pyridazine-3,5-diamine;
 N-{4-[5-(5-Cyclopropyl-1*H*-pyrazol-3-ylamino)-pyridazin-3-ylsulfanyl]-phenyl}-acetamide;
 N⁵-(1*H*-Indazol-3-yl)-N³-(1*H*-indazol-6-yl)-pyridazine-3,5-diamine; and
 (1*H*-Indazol-3-yl)-[6-(3-methoxy-phenylsulfanyl)-pyridazin-4-yl]-amine.

Claim 10 (Original): A composition comprising a compound according to any of claims 1-9, and a pharmaceutically acceptable carrier.

Claim 11 (Original): The composition according to claim 10, further comprising an additional therapeutic agent.

Claim 12 (Original): A method of inhibiting Aurora-2 or GSK-3 activity in a biological sample comprising the step of contacting said biological sample with a compound according to any one of claims 1-9.

Claim 13 (Original): A method of inhibiting Aurora-2 activity in a patient comprising the step of administering to said patient a composition according to claim 10.

Claim 14 (Original): A method of inhibiting Aurora-2 activity in a patient comprising the step of administering to said patient a composition according to claim 11.

Claim 15 (Original): A method of treating an Aurora-2-mediated disease, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a composition according to claim 10.

Claim 16 (Original): The method according to claim 15, wherein said disease is selected from colon, breast, stomach, or ovarian cancer.

Claim 17 (Original): The method according to claim 16, wherein said method further comprises administering an additional therapeutic agent.

Claim 18 (Original): The method according to claim 17, wherein said additional therapeutic agent is a chemotherapeutic agent.

Claim 19 (Original): A method of inhibiting GSK-3 activity in a patient comprising the step of administering to said patient a composition according to claim 10.

Claim 20 (Original): A method of inhibiting GSK-3 activity in a patient comprising the step of administering to said patient a composition according to claim 11.

Claim 21 (Currently amended): A method of ~~method of treating~~ a GSK-3-mediated disease, which method comprises administering to a patient in need of such a treatment a therapeutically effective amount of a composition according to claim 10.

Claim 22 (Original): The method according to claim 21, wherein said GSK-3-mediated disease is selected from diabetes, Alzheimer's disease, Huntington's Disease, Parkinson's Disease, AIDS-associated dementia, amyotrophic lateral sclerosis (AML), multiple sclerosis (MS), schizophrenia, cardiomyocyte hypertrophy, reperfusion/ischemia, or baldness.

Claim 23 (Original): The method according to claim 22, wherein said GSK-3-mediated disease is diabetes.

Claim 24 (Original): A method of enhancing glycogen synthesis or lowering blood levels of glucose in a patient in need thereof, which method comprises administering to said patient a therapeutically effective amount of a composition according to claim 10.

Claim 25 (Original): A method of inhibiting the production of hyperphosphorylated Tau protein in a patient, which method comprises administering to a patient in need thereof a therapeutically effective amount of a composition according to claim 10.

Claim 26 (Original): A method of inhibiting the phosphorylation of β -catenin, which method comprises administering to a patient in need thereof a therapeutically effective amount of a composition according to claim 10.